arXiv:1304.2254v1 [math.NT] 8 Apr 2013

PROOF OF A CONJECTURE ON PERMUTATION POLYNOMIALS OVER FINITE FIELDS

XIANG-DONG HOU*

ABSTRACT. Let k be a positive integer and $S_{2k} = \mathtt{x}+\mathtt{x}^4+\cdots+\mathtt{x}^{4^{2k-1}} \in \mathbb{F}_2[\mathtt{x}]$. It was recently conjectured that $\mathtt{x}+S_{2k}^{4^{2k}}+S_{2k}^{4^k+3}$ is a permutation polynomial of $\mathbb{F}_{4^{3k}}$. In this note, the conjecture is confirmed and a generalization is obtained.

1. INTRODUCTION

Let $\mathbb{F}_q$ denote the finite fields with q elements. For integer $k \ge 0$, define

$$S_{k,q} = \mathtt{x} + \mathtt{x}^q + \cdots + \mathtt{x}^{q^{k-1}} \in \mathbb{F}_p[\mathtt{x}] \qquad (p = \operatorname{char} \mathbb{F}_q).$$

When q is fixed, we write $S_{k,q} = S_k$. The main purpose of this note is to prove the following theorem.

Theorem 1.1. *Let $q = 4$ and $k \ge 1$. Then $\mathtt{x} + S_{2k}^{q^{2k}} + S_{2k}^{q^k+3}$ is a permutation polynomial (PP) of $\mathbb{F}_{q^{3k}}$.*

Theorem 1.1 appeared as a conjecture in a recent study of permutation polynomials over finite fields [1]. We refer the reader to [1] for more background of this conjecture. An adaptation of the proof of Theorem 1.1 gives a slight generalization of Theorem 1.1, which is also included in thos note.

2. PROOF OF THEOREM 1.1

We first recall two facts:

Fact 2.1. [2, Theorem 7.7] Let $f \in \mathbb{F}_q[\mathtt{x}]$. Then f is a PP of $\mathbb{F}_q$ if and only if for all $a \in \mathbb{F}_q^*$, $\sum_{x\in\mathbb{F}_q} \zeta_p^{\operatorname{Tr}_{q/p}(af(x))} = 0$, where $p = \operatorname{char} \mathbb{F}_q$ and $\zeta_p = e^{\frac{2\pi i}{p}}$.

Fact 2.2. [1, Lemma 6.13] Let p be a prime and $f : \mathbb{F}_p^n \to \mathbb{F}_p$ a function. If there exists a $y \in \mathbb{F}_p^n$ such that $f(x+y) - f(x)$ is a nonzero constant for all $x \in \mathbb{F}_p^n$, then $\sum_{x\in\mathbb{F}_p^n} \zeta_p^{f(x)} = 0$.

Proof of Theorem 1.1. Recall that $q = 4$ and $k \ge 1$. Write $g = \mathtt{x} + S_{2k}^{q^{2k}} + S_{2k}^{q^k+3}$ and $\operatorname{Tr} = \operatorname{Tr}_{q^{3k}/2}$. By Fact 2.1, it suffices to show that for every $a \in \mathbb{F}_{q^{3k}}^*$,

$$\sum_{x\in\mathbb{F}_{q^{3k}}} (-1)^{\operatorname{Tr}(ag(x))} = 0. \tag{2.1}$$

2000 *Mathematics Subject Classification.* 11T06, 11T55.

Key words and phrases. exponential sum, finite field, permutation polynomial.

* Research partially supported by NSA Grant H98230-12-1-0245.

We will use the relation

$$S_{2k}+S_{2k}^{q^k}+S_{2k}^{q^{2k}}\equiv 0 \pmod{\mathtt{x}^{q^{3k}}-\mathtt{x}}. \tag{2.2}$$

Case 1. Assume $\mathrm{Tr}_{q^{3k}/q^k}(a)\neq 0$. Then there exists a $y\in\mathbb{F}_{q^k}$ such that $\mathrm{Tr}_{q^k/2}\bigl[y\mathrm{Tr}_{q^{3k}/q^k}(a)\bigr]\neq 0$. For all $x\in\mathbb{F}_{q^{3k}}$ we have $S_{2k}(x+y)=S_{2k}(x)+S_{2k}(y)=S_{2k}(x)$. Hence

$$\mathrm{Tr}\bigl[ag(x+y)-ag(x)\bigr]=\mathrm{Tr}(ay)=\mathrm{Tr}_{q^k/2}\bigl[y\mathrm{Tr}_{q^{3k}/q^k}(a)\bigr],$$

which is a nonzero constant. By Fact 2.2, (2.1) holds.

Case 2. Assume $\mathrm{Tr}_{q^{3k}/q^k}(a)=0$. The $a=c+c^{q^k}$ for some $c\in\mathbb{F}_{q^{3k}}\setminus\mathbb{F}_{q^k}$. For $x\in\mathbb{F}_{q^{3k}}$, we write $S_{2k}(x)=S_{2k}$ (a slight abuse of notation). We have

$$\begin{aligned}
&\mathrm{Tr}\bigl(ag(x)\bigr)\\
&=\mathrm{Tr}\bigl[(c+c^{q^k})g(x)\bigr]\\
&=\mathrm{Tr}\bigl[c\bigl(g(x)+g(x)^{q^{2k}}\bigr)\bigr]\\
&=\mathrm{Tr}\bigl[c(x+S_{2k}^{q^{2k}}+S_{2k}^{q^k+3}+x^{q^{2k}}+S_{2k}^{q^k}+S_{2k}^{1+3q^{2k}})\bigr]\\
&=\mathrm{Tr}\bigl[c(x+x^{q^{2k}}+S_{2k}+S_{2k}^3(S_{2k}+S_{2k}^{q^{2k}})+S_{2k}^{1+3q^{2k}})\bigr] && \text{(by (2.2))}\\
&=\mathrm{Tr}\bigl[c(S_{2k}^q+S_{2k}^4+S_{2k}^{3+q^{2k}}+S_{2k}^{1+3q^{2k}})\bigr]\\
&=\mathrm{Tr}\bigl[cS_{2k}^{1+q^{2k}}(S_{2k}^2+S_{2k}^{2q^{2k}})\bigr]\\
&=\mathrm{Tr}\bigl[cS_{2k}^{1+q^{2k}}(S_{2k}+S_{2k}^{q^{2k}})^2\bigr]\\
&=\mathrm{Tr}(cS_{2k}^{1+q^{2k}}S_{2k}^{2q^k}) && \text{(by (2.2))}\\
&=\mathrm{Tr}(cS_{2k}^{1+2q^k+q^{2k}}).
\end{aligned} \tag{2.3}$$

By (2.1), $S_{2k}(\mathbb{F}_{q^{2k}})\subset\mathrm{Tr}_{q^{3k}/q^k}^{-1}(0)$. Since

$$\gcd(1+\mathtt{x}+\cdots+\mathtt{x}^{2k-1},\mathtt{x}^{3k}-1)=\mathtt{x}^k-1,$$

by [2, Theorem 3.62], $\{x\in\mathbb{F}_{q^{3k}}:S_{2k}(x)=0\}=\mathbb{F}_{q^k}$. Thus $S_{2k}:\mathbb{F}_{q^{3k}}\to\mathrm{Tr}_{q^{3k}/q^k}^{-1}(0)$ is an onto $\mathbb{F}_{q^k}$-map with $\ker S_{2k}=\mathbb{F}_{q^k}$. Therefore

$$\begin{aligned}
\sum_{x\in\mathbb{F}_{q^{3k}}}(-1)^{\mathrm{Tr}(ag(x))}&=\sum_{x\in\mathbb{F}_{q^{3k}}}(-1)^{\mathrm{Tr}(cS_{2k}^{1+2q^k+q^{2k}})} && \text{(by (2.3))}\\
&=q^k\sum_{x\in\mathrm{Tr}_{q^{3k}/q^k}^{-1}(0)}(-1)^{\mathrm{Tr}(cx^{1+2q^k+q^{2k}})}.
\end{aligned} \tag{2.4}$$

Let d_1, d_2 be a basis of $\mathrm{Tr}^{-1}_{q^{3k}/q^k}(0)$ over $\mathbb{F}_{q^k}$. We have

$$\begin{aligned}
&\sum_{x\in\mathrm{Tr}^{-1}_{q^{3k}/q^k}(0)} (-1)^{\mathrm{Tr}(cx^{1+2q^k+q^{2k}})}\\
&= \sum_{x,y\in\mathbb{F}_{q^k}} (-1)^{\mathrm{Tr}[c(d_1x+d_2y)^{1+2q^k+q^{2k}}]}\\
&= \sum_{x\in\mathbb{F}_{q^k}} (-1)^{\mathrm{Tr}[c(d_1x)^{1+2q^k+q^{2k}}]} + \sum_{\substack{x\in\mathbb{F}_{q^k}\\ y\in\mathbb{F}^*_{q^k}}} (-1)^{\mathrm{Tr}[c(d_1x+d_2y)^{1+2q^k+q^{2k}}]}\\
&= \sum_{x\in\mathbb{F}_{q^k}} (-1)^{\mathrm{Tr}(cd_1^{q^k}x)} + \sum_{\substack{x\in\mathbb{F}_{q^k}\\ y\in\mathbb{F}^*_{q^k}}} (-1)^{\mathrm{Tr}[c(d_1xy+d_2y)^{1+2q^k+q^{2k}}]}
\end{aligned}$$

(for the first sum, $x \leftarrow d_1^{1+q^k+q^{2k}} x^{1+2q^k+q^{2k}}$; for the second sum $x \to xy$)

$$= \sum_{x\in\mathbb{F}_{q^k}} (-1)^{\mathrm{Tr}(cd_1^{q^k}x)} + \sum_{\substack{x\in\mathbb{F}_{q^k}\\ y\in\mathbb{F}^*_{q^k}}} (-1)^{\mathrm{Tr}[c(d_1x+d_2)^{q^k}y]} \tag{2.5}$$

(for the first sum, $y \leftarrow (d_1x+d_2)^{1+q^k+q^{2k}} y^{1+2q^k+q^{2k}}$)

$$\begin{aligned}
&= \sum_{x\in\mathbb{F}_{q^k}} (-1)^{\mathrm{Tr}(cd_1^{q^k}x)} + \sum_{\substack{x\in\mathbb{F}_{q^k}\\ y\in\mathbb{F}^*_{q^k}}} (-1)^{\mathrm{Tr}(cd_1^{q^k}xy+cd_2^{q^k}y)}\\
&= \sum_{x\in\mathbb{F}_{q^k}} (-1)^{\mathrm{Tr}(cd_1^{q^k}x)} + \sum_{\substack{x\in\mathbb{F}_{q^k}\\ y\in\mathbb{F}^*_{q^k}}} (-1)^{\mathrm{Tr}(cd_1^{q^k}x+cd_2^{q^k}y)}
\end{aligned}$$

(for the second sum, $x \leftarrow xy$)

$$\begin{aligned}
&= \Big[\sum_{x\in\mathbb{F}_{q^k}} (-1)^{\mathrm{Tr}(cd_1^{q^k}x)}\Big]\Big[1+\sum_{y\in\mathbb{F}^*_{q^k}} (-1)^{\mathrm{Tr}(cd_2^{q^k}y)}\Big]\\
&= \Big[\sum_{x\in\mathbb{F}_{q^k}} (-1)^{\mathrm{Tr}(cd_1^{q^k}x)}\Big]\Big[\sum_{y\in\mathbb{F}_{q^k}} (-1)^{\mathrm{Tr}(cd_2^{q^k}y)}\Big].
\end{aligned}$$

We claim that $\mathrm{Tr}_{q^{3k}/q^k}(cd_1^{q^k})$ and $\mathrm{Tr}_{q^{3k}/q^k}(cd_2^{q^k})$ cannot be both 0. (Otherwise, $\mathrm{Tr}^{-1}_{q^{3k}/q^k}(0) = d_1^{q^k}\mathbb{F}_{q^k} + d_2^{q^k}\mathbb{F}_{q^k}$ would be a vector space over $\mathbb{F}_{q^k}(c) = \mathbb{F}_{q^{3k}}$, which is impossible.) Therefore, at least one of the two sums in the last line of (2.3) is 0, so we have

$$\sum_{x\in\mathrm{Tr}^{-1}_{q^{3k}/q^k}(0)} (-1)^{\mathrm{Tr}(cx^{1+2q^k+q^{2k}})} = 0. \tag{2.6}$$

Combining (2.4) and (2.6) gives (2.1). □

3. Generalization

The proof in Section 3 works for the following generalization of Theorem 1.1

Theorem 3.1. *Let q be a power of 2. Let $L \in \mathbb{F}_{q^{3k}}[\mathtt{x}]$ be a 2-linearized polynomial such that*

- (i) *L permutes $\mathbb{F}_{q^k}$, and*
- (ii) *$L + L^{q^{2k}} \equiv S_{2k}^4 \pmod{\mathtt{x}^{q^{3k}} - \mathtt{x}}$.*

Then $L + S_{2k}^{q^k+3}$ is a PP of $\mathbb{F}_{q^{3k}}$.

Note. $L = (\mathtt{x} + S_{2k}^{q^{2k}})^{4q^{3k-1}}$ satisfies conditions (i) and (ii) of Theorem 3.1. (i) is obvious. For (ii), we have $L + L^{q^{2k}} = (\mathtt{x} + S_{2k}^{q^{2k}} + \mathtt{x}^{q^{2k}} + S_{2k}^{q^k})^{4q^{3k-1}} \equiv (\mathtt{x} + \mathtt{x}^{q^{2k}} + S_{2k})^{4q^{3k-1}} = (S_{2k}^q)^{4q^{3k-1}} \equiv S_{2k}^4$, where "$\equiv$" means "$\equiv \pmod{\mathtt{x}^{q^{3k}} - \mathtt{x}}$".

Proof of Theorem 3.1. Let $g = L + S_{2k}^{q^k+3}$ and follow the proof of Theorem 1.1. In Case 1, we choose $y \in \mathbb{F}_{q^k}$ such that $\mathrm{Tr}_{q^k/2}\big[L(y)\mathrm{Tr}_{q^{3k}/q^k}(a)\big] \neq 0$. In Case 2, (2.3), still holds because of condition (ii). $\square$

REFERENCES

[1] N. Fernando, X. Hou, S. D. Lappano, *A new approach to permutation polynomials over finite fields, II*, Finite Fields Appl., Available online 23 January 2013

[2] R. Lidl and H. Niederreiter, *Finite Fields*, 2nd ed., Cambridge Univ. Press, Cambridge, 1997.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SOUTH FLORIDA, TAMPA, FL 33620

E-mail address: `xhou@usf.edu`